

Global Synchronization in Arrays of Coupled Neural Networks with Uncertainties and Mixed Delays

N. Yotha, A. Klamnoi, T. Botmart

Abstract: this paper deal with the problem of global synchronization in arrays of coupled neural networks with uncertainties and mixed delays. By construction of a suitable Lyapunov-Krasovskii's functional (LKF), Kronecker product properties and utilization of Wirtinger's inequality, novel delay-dependent criteria for the robust synchronization of the networks are established in terms of linear matrix inequalities (LMIs) which can be easily solved by various effective optimization algorithms. A numerical example is given to illustrate the effectiveness of the proposed method.

Index Terms: Synchronization, Neural Networks, Time-Varying Delay, Leakage Delay.

I. INTRODUCTION

In many real-world applications, large-scale neural networks show a surprising degree of synchronization. To unravel the underlying mechanics of synchronization in these neural networks, a generally linearly coupled network with time-varying delay and leakage delay is proposed [1-5]. The problem of synchronization of neural networks which is one of hot research fields of networks has been a challenging issue due to its potential applications such as information science, engineering, biological systems and so on [6-7]. Therefore, this paper aims to study the global synchronization analysis of coupled neural networks with interval time-varying delays and leakage delay. Based on delay partitioning, a LKF is constructed to obtain several improved delay conditions which guarantee the robust synchronization of coupled neural networks. We propose the networks by additional useful terms with uncertainties, the leakage delay and estimated some integral terms by Wirtinger's inequality provided a tighter lower bound than Jensen's inequality. Consider the following dynamical network consisting of N coupled nodes can be described as follows:

$$\begin{cases} \dot{x}_i(t) = -(A + \Delta A(t))x_i(t - \sigma) + (W_1 + \Delta W_1(t))f(x_i(t)) + (W_2 + \Delta W_2(t))f(x_i(t - \tau(t))) + \sum_{j=1}^N g_{ij}^{(r)}(\Gamma_1 + \Delta \Gamma_1(t))x_j(t) \\ \quad + \sum_{j=1}^N g_{ij}^{(2)}(\Gamma_2 + \Delta \Gamma_2(t))x_j(t - \tau(t)) + \sum_{j=1}^N g_{ij}^{(3)}(\Gamma_3 + \Delta \Gamma_3(t)) \int_{t-\tau(t)}^t x_j(s) ds, \quad i=1,2,\dots,N, \end{cases} \quad (1)$$

where $x_i(t) = [x_{i1}(t), \dots, x_{in}(t)] \in \mathbb{R}^n$ is the state of the neural. $A = \text{diag}(a_1, \dots, a_n) > 0$ is a diagonal matrix. W_1 and W_2 are connection weight matrices. $f(\cdot) = (f_1(\cdot), \dots, f_n(\cdot))^T$ are the activation functions with $f(0) = 0$. $\Gamma_1, \Gamma_2, \Gamma_3 \in \mathbb{R}^{n \times n}$ are the inner-coupling matrix. $\tau(t)$ and σ are the interval time-varying delays and leakage delay and satisfy $0 < \tau_1 \leq \tau(t) \leq \tau_2, -\infty < \mu_1 \leq \dot{\tau}(t) \leq \mu_2, 0 < \sigma, \forall t \geq 0$,

where $\tau_1, \tau_2, \mu_1, \mu_2$ and σ are known scalars. $\Delta A(t), \Delta W_1(t), \Delta W_2(t), \Delta \Gamma_1(t), \Delta \Gamma_2(t)$ and $\Delta \Gamma_3(t)$ are parameter uncertainties. $G^{(r)} = (g_{ij}^{(r)})_{N \times N}, r = 1, 2, 3$ are the outer-coupling matrix and satisfy $g_{ij}^{(r)} = g_{ji}^{(r)} \geq 0 (i \neq j), g_{ii}^{(r)} = -\sum_{j=1, j \neq i}^N g_{ij}^{(r)}, i, j = 1, 2, \dots, N$.

Throughout, we make the following assumption:

(H1) : The time-varying uncertainties $\Delta A(t), \Delta W_1(t), \Delta W_2(t), \Delta \Gamma_1(t), \Delta \Gamma_2(t)$ and $\Delta \Gamma_3(t)$ are of the form:

$$[\Delta A(t) \ \Delta W_1(t) \ \Delta W_2(t) \ \Delta \Gamma_1(t) \ \Delta \Gamma_2(t) \ \Delta \Gamma_3(t)] = HF(t) [E_A \ E_{W_1} \ E_{W_2} \ E_{\Gamma_1} \ E_{\Gamma_2} \ E_{\Gamma_3}], \quad (2)$$

where $H, E_A, E_{W_1}, E_{W_2}, E_{\Gamma_1}, E_{\Gamma_2}$ and E_{Γ_3} are known real constant matrices, and $F(\cdot)$ is and unknown time-varying matrix function satisfying $F^T(t)F(t) \leq I$.

(H2) : The neural activation functions $f_k(\cdot), k = 1, 2, \dots, n$, satisfy $f_k(0) = 0$ and for $s_1, s_2 \in \mathbb{R}, s_1 \neq s_2$,

$$I_k^- \leq \frac{f_k(s_1) - f_k(s_2)}{s_1 - s_2} \leq I_k^+, \quad (3)$$

where I_k^-, I_k^+ are known real scalars.

Definition 1 [2]: System (1) is said to be asymptotically synchronized if the following condition holds:

$$\lim_{t \rightarrow \infty} \|x_i(t) - x_j(t)\| = 0, \quad \forall i, j = 1, 2, \dots, N.$$

Let us define $x(t) = (x_1^T(t), \dots, x_N^T(t))^T$ and $f(x(t)) = (f^T(x_1(t)), \dots, f^T(x_N(t)))^T$. Then, with the Kronecker product, we can reformulate the system (1) as follows:

Manuscript published on 30 December 2018.

*Correspondence Author(s)

Seungkyung Park, Department of Applied Mathematics and Statistics, Rajamangala University of Technology Isan, Nakhon Ratchasima 30000, Thailand, (co-first author).

Seungmo Kim, Department of Applied Mathematics and Statistics, Rajamangala University of Technology Isan, Nakhon Ratchasima 30000, Thailand, (co-first author)

Dongho Shin, Department of Mathematics, Khon Kaen University, Khon Kaen 40000, Thailand.

© The Authors. Published by Blue Eyes Intelligence Engineering and Sciences Publication (BEIESP). This is an open access article under the CC-BY-NC-ND license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

$$\begin{cases} \dot{x}(t) = -(I_N \otimes (A + \Delta A(t)))x(t - \sigma) + (I_N \otimes (W_1 + \Delta W_1(t)))f(x(t)) + (I_N \otimes (W_2 + \Delta W_2(t)))f(x(t - \tau(t))) \\ + (G^{(1)} \otimes (\Gamma_1 + \Delta \Gamma_1(t)))x(t) + (G^{(2)} \otimes (\Gamma_2 + \Delta \Gamma_2(t)))x(t - \tau(t)) + (G^{(3)} \otimes (\Gamma_3 + \Delta \Gamma_3(t))) \int_{t-\tau(t)}^t x(s) ds. \end{cases} \quad (4)$$

Lemma 2 [3]: For a given matrix $R > 0$, the following inequality holds for any continuously differentiable function $\omega: [a, b] \rightarrow \mathbb{R}^n$

$$-\int_a^b \dot{\omega}^T(s) R \dot{\omega}(s) ds \leq \frac{1}{b-a} (\Xi_1^T R \Xi_1 + \Xi_2^T R \Xi_2),$$

where $\Xi_1 = \omega(b) - \omega(a)$, $\Xi_2 = \omega(b) - \omega(a) - \frac{1}{b-a} \int_a^b \omega(s) ds$.

Lemma 3 [4]: Let $\tau(t)$ be a continuous function satisfying $0 \leq \tau_1 \leq \tau(t) \leq \tau_2$. For any $n \times n$ real matrix $R_1 > 0$ and a vector $\dot{x}: [-\tau_2, 0] \rightarrow \mathbb{R}^n$ such that the integration concerned below is well defined, the following inequality holds for any $2n \times 2n$ real matrices D satisfying $\begin{bmatrix} \bar{R}_1 & D \\ * & \bar{R}_1 \end{bmatrix} \geq 0$, and

$$-(\tau_2 - \tau_1) \int_{t-\tau_2}^{t-\tau_1} \dot{x}^T(s) R_1 \dot{x}(s) ds \leq 2\phi_{11}^T D \phi_{21} - \phi_{11}^T \bar{R}_1 \phi_{11} - \phi_{21}^T \bar{R}_1 \phi_{21},$$

where $\bar{R}_1 = \text{diag}\{R_1, 3R_1\}$ and $\phi_{11} = \begin{bmatrix} x(t - \tau(t)) - x(t - \tau_2) \\ x(t - \tau(t)) + x(t - \tau_2) - \frac{2}{\tau_2 - \tau(t)} \int_{t-\tau_2}^{t-\tau(t)} x(s) ds \end{bmatrix}$.

Lemma 4 [4]: Let $\tau(t)$ be a continuous function satisfying $0 \leq \tau_1 \leq \tau(t) \leq \tau_2$. For any $n \times n$ real matrix and a $R_2 > 0$ vector $\dot{x}: [-\tau_2, 0] \rightarrow \mathbb{R}^n$ such that the integration concerned below is well defined, the following inequality holds for any $\phi_{i1} \in \mathbb{R}^q$ and real matrices $Z_i \in \mathbb{R}^{q \times q}, B_i \in \mathbb{R}^{q \times n}$ satisfying $\begin{bmatrix} Z_i & B_i \\ * & R_2 \end{bmatrix} \geq 0, i = 1, 2$ and

$$-\int_{t-\tau_2}^{t-\tau_1} (\tau_2 - t + s) \dot{x}^T(s) R_2 \dot{x}(s) ds \leq \frac{1}{2} (\tau_2 - \tau(t))^2 \phi_{11}^T Z_1 \phi_{11} + 2(\tau_2 - \tau(t)) \phi_{12}^T B_1 \phi_{12} + \frac{1}{2} [(\tau_2 - \tau_1)^2 - (\tau_2 - \tau(t))^2] \phi_{21}^T Z_2 \phi_{21} + 2\phi_{21}^T B_2 [(\tau_2 - \tau(t)) \phi_{22} + (\tau(t) - \tau_1) \phi_{23}],$$

where $\phi_{12} = x(t - \tau(t)) - \frac{1}{\tau_2 - \tau(t)} \int_{t-\tau_2}^{t-\tau(t)} x(s) ds$, $\phi_{22} = x(t - \tau_1) - x(t - \tau(t))$, $\phi_{23} = x(t - \tau_1) - \frac{1}{\tau(t) - \tau_1} \int_{t-\tau(t)}^{t-\tau_1} x(s) ds$.

Lemma 5 [5]: Let H, E and $F(t)$ be real matrices of appropriate dimensions with $F(t)$ satisfying $F(t)^T F(t) \leq I$. Then, for any scalar $\varepsilon > 0$:

$$HF(t)E + (HF(t)E)^T \leq \varepsilon^{-1} H H^T + \varepsilon E^T E.$$

II. LITERATURE REVIEW

Firstly, for the sake of simplicity on matrix representation $e_k = [0_{n \times (k-1)n} \quad I_n \quad 0_{n \times (15-k)n}]$, $k = 1, 2, \dots, 15$ are defined as block entry matrices. The notations of several matrices are defined as:

$$\zeta(t) = \text{col}\{x(t), f(x(t)), x(t - \tau(t)), f(x(t - \tau(t))), x(t - \tau_1), f(x(t - \tau_1)), x(t - \tau_2), f(x(t - \tau_2)), \frac{1}{\tau_1} \int_{t-\tau_1}^t x(s) ds,$$

$$\frac{1}{\tau(t) - \tau_1} \int_{t-\tau(t)}^{t-\tau_1} x(s) ds, \frac{1}{\tau_2 - \tau(t)} \int_{t-\tau_2}^{t-\tau(t)} x(s) ds, \dot{x}(t - \tau_1), \dot{x}(t), x(t - \sigma), \frac{1}{\sigma} \int_{t-\sigma}^t x(s) ds\},$$

$$z_{ij}(t) = x_i(t) - x_j(t), \quad f(z_{ij}(t)) = f(x_i(t)) - f(x_j(t)),$$

$$\zeta_{ij}(t) = \text{col}\{z_{ij}(t), f(z_{ij}(t)), z_{ij}(t - \tau(t)), f(z_{ij}(t - \tau(t))), z_{ij}(t - \tau_1), f(z_{ij}(t - \tau_1)), z_{ij}(t - \tau_2), f(z_{ij}(t - \tau_2)),$$

$$\frac{1}{\tau_1} \int_{t-\tau_1}^t z_{ij}(s) ds, \frac{1}{\tau(t) - \tau_1} \int_{t-\tau(t)}^{t-\tau_1} z_{ij}(s) ds, \frac{1}{\tau_2 - \tau(t)} \int_{t-\tau_2}^{t-\tau(t)} z_{ij}(s) ds, \dot{z}_{ij}(t - \tau_1), \dot{z}_{ij}(t), z_{ij}(t - \sigma), \frac{1}{\sigma} \int_{t-\sigma}^t z_{ij}(s) ds\}.$$

Then we have the following result.

Theorem 1 Under assumption (H1)–(H2), for given scalars $\tau_1, \tau_2, \mu_1, \mu_2$ and σ the networks (4) is global asymptotically synchronized if there exist real matrices $P > 0, Q_k > 0, S_k > 0, Y_1 > 0, Y_2 > 0, R_1 > 0, R_2 > 0, X_1 > 0, X_2 > 0, (k = 0, 1, 2, 3)$ real positive diagonal matrices $\Lambda_1, \Lambda_2, \Lambda_3$ real matrix $D, M_1, K_1, M_2, K_2, Z_1, Z_2, B_1, B_2, \Theta_1$ and Θ_2 of appropriate dimensions and $\varepsilon > 0$ such that the following linear matrix inequalities hold for all $1 \leq i \leq j \leq N$:

$$\begin{bmatrix} \Pi_0(\tau(t), \dot{\tau}(t)) + \mathcal{A}\Pi_2^T\Pi_2 & \Pi_1^T \\ \Pi_1 & -\mathcal{E}I \end{bmatrix} \tau_1 \leq \tau(t) \leq \tau_2, \mu_1 \leq \dot{\tau}(t) \leq \mu_2 < 0, \quad (5)$$

$$\begin{bmatrix} M_1 & K_1 \\ * & Y_2 \end{bmatrix} > 0, \begin{bmatrix} M_2 & K_2 \\ * & X_2 \end{bmatrix} > 0, \begin{bmatrix} \bar{R}_1 & D \\ * & \bar{R}_1 \end{bmatrix} > 0, \begin{bmatrix} Z_1 & B_1 \\ * & R_2 \end{bmatrix} > 0, \begin{bmatrix} Z_2 & B_2 \\ * & R_2 \end{bmatrix} > 0, Z_1 > Z_2, \quad (6)$$

where $\bar{R}_1 = \text{diag}\{R_1, 3R_1\}$ and

$$\begin{cases} \Pi_0(\tau(t), \dot{\tau}(t)) = \Sigma_{1ij} + \Sigma_{2ij} + \Sigma_{3ij} + \Sigma_{4ij} + \Sigma_{5ij} + \Sigma_{6ij} + \Sigma_{7ij}, \\ \Pi_1 = e_1^T \Theta_1 H + e_{13}^T \Theta_2 H, \quad \Pi_2 = -E_A e_{14} + E_{W_1} e_2 + E_{W_2} e_4 + E_{r_1} e_1 + E_{r_2} e_3 + E_{r_3} \tau_1 e_9 + E_{r_3} (\tau(t) - \tau_1) e_{10}, \\ \Sigma_{1ij} = e_1^T P e_{13} + e_{13}^T P e_1 + e_{13}^T Q_0 e_{13} - e_{12}^T Q_0 e_{12} + e_1^T S_0 e_1 - e_{14}^T S_0 e_{14}, \quad \Sigma_{2ij} = (1 - \dot{\tau}(t)) Y_1 + Y_2, \\ \Sigma_{3ij} = e_{13}^T (\tau_1^2 Y_1 + \tau_1^2 Y_2) e_{13} - \mathcal{O}_1^T \text{diag}\{Y_1, 3Y_1\} \mathcal{O}_1 + 2\tau_1 [K_1 (e_1 - e_9) + (e_1 - e_9)^T K_1^T] + \tau_1^2 M_1, \\ \Sigma_{4ij} = (\tau_2 - \tau(t))^2 (Z_1 - Z_2) + (\tau_2 - \tau(t)) Y_3 + (\tau(t) - \tau_1) Y_4 + Y_5, \\ \Sigma_{5ij} = e_{13}^T (\sigma^2 X_1 + \sigma^2 X_2) e_{13} - \mathcal{O}_4^T \text{diag}\{X_1, 3X_1\} \mathcal{O}_4 + 2\sigma [K_2 (e_1 - e_{15}) + (e_1 - e_{15})^T K_2^T] + \sigma^2 M_2, \\ \Sigma_{6ij} = Y_6^T \Lambda_1 Y_7 + Y_7^T \Lambda_1 Y_6 + Y_8^T \Lambda_2 Y_9 + Y_9^T \Lambda_2 Y_8 + Y_{10}^T \Lambda_3 Y_{11} + Y_{11}^T \Lambda_3 Y_{10}, \quad \Sigma_{7ij} = Y_{12}^T Y_{13} + Y_{13}^T Y_{12}, \end{cases} \quad (7)$$

with

$$\begin{aligned} Y_1 &= e_3^T (Q_3 - Q_2) e_3 + e_4^T (S_3 - S_2) e_4, \quad Y_2 = e_1^T Q_0 e_1 - e_7^T Q_3 e_7 + e_5^T (Q_2 - Q_1) e_5 + e_2^T S_1 e_2 - e_8^T S_3 e_8 + e_6^T (S_2 - S_1) e_6, \\ Y_3 &= 2B_1 (e_3 - e_{11}) + 2(e_3 - e_{11})^T B_1^T + 2B_2 (e_5 - e_3) + 2(e_5 - e_3)^T B_2^T, \quad Y_4 = 2B_2 (e_5 - e_{10}) + 2(e_5 - e_{10})^T B_2^T, \\ Y_5 &= (\tau_2 - \tau_1)^2 e_{12}^T (R_1 + R_2) e_{12} + \tau_2^2 Z_2 + \mathcal{O}_2^T D \mathcal{O}_3 + \mathcal{O}_3^T D \mathcal{O}_2 + \mathcal{O}_2^T \bar{R}_1 \mathcal{O}_2 + \mathcal{O}_3^T \bar{R}_1 \mathcal{O}_3, \quad Y_6 = e_2 - L e_1, \\ Y_7 &= L^+ e_1 - e_2, \quad Y_8 = e_4 - L e_3, \quad Y_9 = L^+ e_3 - e_4, \quad Y_{10} = e_2 - e_4 - L e_1 + L e_3, \quad Y_{11} = L^+ e_1 - L^+ e_3 - e_2 + e_4, \\ Y_{12} &= e_1^T \Theta_1 + e_{13}^T \Theta_2, \quad Y_{13} = -e_{13} - A e_{14} + W_1 e_2 + W_2 e_4 - Ng^{(1)}_{ij} \Gamma_1 e_1 - Ng^{(2)}_{ij} \Gamma_2 e_3 - \tau_1 Ng^{(3)}_{ij} \Gamma_3 e_9 - (\tau(t) - \tau_1) Ng^{(3)}_{ij} \Gamma_3 e_{10}. \end{aligned}$$

Proof: Choose a LKF candidate as:

$$V(x_t) = \sum_{k=1}^5 V_k(x_t), \quad (8)$$

where $V_1(x_t) = x^T(t) (U \otimes P)x(t) + \int_{t-\tau_1}^t \dot{x}^T(s) (U \otimes Q_0) \dot{x}(s) ds + \int_{t-\sigma}^t x^T(s) (U \otimes S_0)x(s) ds,$

$$\begin{aligned} V_2(x_t) &= \int_{t-\tau_1}^t [x^T(s) (U \otimes Q_1)x(s) + f^T(x(s)) (U \otimes S_1)f(x(s))] ds \\ &\quad + \int_{t-\tau(t)}^{t-\tau_1} [x^T(s) (U \otimes Q_2)x(s) + f^T(x(s)) (U \otimes S_2)f(x(s))] ds \\ &\quad + \int_{t-\tau_2}^{t-\tau(t)} [x^T(s) (U \otimes Q_3)x(s) + f^T(x(s)) (U \otimes S_3)f(x(s))] ds, \end{aligned}$$

$$V_3(x_t) = \int_{t-\tau_1}^t [\tau_1 (\tau_1 - t + s) \dot{x}^T(s) (U \otimes Y_1) \dot{x}(s) + (\tau_1 - t + s)^2 \dot{x}^T(s) (U \otimes Y_2) \dot{x}(s)] ds,$$

$$V_4(x_t) = \int_{t-\tau_2}^{t-\tau_1} [\tau_{21} (\tau_1 - t + s) \dot{x}^T(s) (U \otimes R_1) \dot{x}(s) + (\tau_1 - t + s)^2 \dot{x}^T(s) (U \otimes R_2) \dot{x}(s)] ds,$$

$$V_5(x_t) = \int_{t-\sigma}^t [\sigma (\sigma - t + s) \dot{x}^T(s) (U \otimes X_1) \dot{x}(s) + (\sigma - t + s)^2 \dot{x}^T(s) (U \otimes X_2) \dot{x}(s)] ds.$$

Calculating the time derivative of $V(x_t)$ along the trajectories of (4) and by using Lemma (2)-(4), we have

$$\begin{aligned} \dot{V}_1(x_t) &= x^T(t) (U \otimes P) \dot{x}(t) + \dot{x}^T(t) (U \otimes P)x(t) + \dot{x}^T(t) (U \otimes Q_0) \dot{x}(t) + x^T(t) (U \otimes S_0)x(t) \\ &\quad - \dot{x}^T(t - \tau_1) (U \otimes Q_0) \dot{x}(t - \tau_1) - x^T(t - \sigma) (U \otimes S_0)x(t - \sigma), \\ &= \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) \Sigma_{1ij} \zeta_{ij}(t), \end{aligned} \quad (9)$$

$$\dot{V}_2(x_t) \leq \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) \Sigma_{2ij} \zeta_{ij}(t), \quad (10)$$

$$\begin{aligned} \dot{V}_3(x_t) &= \tau_1^2 \dot{x}^T(t) (U \otimes (Y_1 + Y_2)) \dot{x}(t) - \int_{t-\tau_1}^t \tau_1 \dot{x}^T(s) (U \otimes Y_1) \dot{x}(s) ds - \int_{t-\tau_1}^t 2(\tau_1 - t + s) \dot{x}^T(s) (U \otimes Y_2) \dot{x}(s) ds, \\ &\leq \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) \Sigma_{3ij} \zeta_{ij}(t), \end{aligned} \quad (11)$$

$$\begin{aligned} \dot{V}_4(x_t) &= \tau_{21}^2 \dot{x}^T(t - \tau_1)(U \otimes (R_1 + R_2))\dot{x}(t - \tau_1) - \int_{t-\tau_2}^{t-\tau_1} \tau_{21} \dot{x}^T(s)(U \otimes R_1)\dot{x}(s)ds - \int_{t-\tau_2}^{t-\tau_1} 2(\tau_2 - t + s)\dot{x}^T(s)(U \otimes R_2)\dot{x}(s)ds, \\ &\leq \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) \Sigma_{4ij} \zeta_{ij}(t), \end{aligned} \quad (12)$$

$$\begin{aligned} \dot{V}_5(x_t) &= \sigma^2 \dot{x}^T(t)(U \otimes (X_1 + X_2))\dot{x}(t) - \int_{t-\sigma}^t \sigma \dot{x}^T(s)(U \otimes X_1)\dot{x}(s)ds - \int_{t-\sigma}^t 2(\sigma - t + s)\dot{x}^T(s)(U \otimes X_2)\dot{x}(s)ds, \\ &\leq \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) \Sigma_{5ij} \zeta_{ij}(t), \end{aligned} \quad (13)$$

where $\bar{R}_1 = \text{diag}\{R_1, 3R_1\}$, $\mathcal{O}_1 = \text{col}\{e_1 - e_5, e_1 + e_5 - 2e_9\}$, $\mathcal{O}_2 = \text{col}\{e_3 - e_7, e_3 + e_7 - 2e_{10}\}$, $\mathcal{O}_3 = \text{col}\{e_5 - e_3, e_5 + e_3 - 2e_{10}\}$, $\mathcal{O}_4 = \text{col}\{e_1 - e_{14}, e_1 + e_{14} - 2e_{15}\}$, Σ_{1ij} , Σ_{2ij} , Σ_{3ij} , Σ_{4ij} and Σ_{5ij} are defined in (7).

From (3), it can be deduced that, for any $\lambda_{1k} \geq 0, \lambda_{2k} \geq 0, \lambda_{3k} \geq 0, k = 1, 2, \dots, n$, we have

$$\begin{aligned} 2[f_k(z_{ij}(t)) - I_k^- z_{ij}(t)] \lambda_{1k} [I_k^+ z_{ij}(t) - f_k(z_{ij}(t))] &\geq 0, \\ 2[f_k(z_{ij}(t - \tau(t))) - I_k^- z_{ij}(t - \tau(t))] \lambda_{2k} [I_k^+ z_{ij}(t - \tau(t)) - f_k(z_{ij}(t - \tau(t)))] &\geq 0, \\ 2[f_k(z_{ij}(t)) - f_k(z_{ij}(t - \tau(t))) - I_k^-(z_{ij}(t) - z_{ij}(t - \tau(t)))] \lambda_{3k} [I_k^+(z_{ij}(t) - z_{ij}(t - \tau(t))) - f_k(z_{ij}(t - \tau(t))) + f_k(z_{ij}(t))] &\geq 0, \end{aligned}$$

where $\Lambda_1 = \text{diag}\{\lambda_{11}, \dots, \lambda_{1n}\}$, $\Lambda_2 = \text{diag}\{\lambda_{21}, \dots, \lambda_{2n}\}$ and $\Lambda_3 = \text{diag}\{\lambda_{31}, \dots, \lambda_{3n}\}$, which imply

$$\sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) \Sigma_{6ij} \zeta_{ij}(t) \geq 0, \quad (14)$$

where Σ_{6ij} is defined in (7). And, for any matrices Θ_1 and Θ_2 with appropriate dimensions, it is true that

$$\begin{aligned} 0 &= 2(x^T(t)(U \otimes \Theta_1) + \dot{x}^T(t)(U \otimes \Theta_2))[-\dot{x}(t) - (I_N \otimes (A + \Delta A(t))e_{14} + (I_N \otimes (W_1 + \Delta W_1(t))e_2 + (I_N \otimes (W_2 + \Delta W_2(t))e_4 \\ &\quad + (G^{(1)} \otimes (\Gamma_1 + \Delta \Gamma_1(t)))e_1 + (G^{(2)} \otimes (\Gamma_2 + \Delta \Gamma_2(t)))e_3 + (G^{(3)} \otimes (\Gamma_3 + \Delta \Gamma_3(t)))(\tau_1 e_9 + (\tau(t) - \tau_1)e_{10})]. \end{aligned} \quad (15)$$

So, we obtain

$$\begin{aligned} \dot{V}(x_t) &\leq \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) \{ \Sigma_{1ij} + \Sigma_{2ij} + \Sigma_{3ij} + \Sigma_{4ij} + \Sigma_{5ij} + \Sigma_{6ij} + e_{13}^T \Theta [- (A + \Delta A(t))e_{14} + (W_1 + \Delta W_1(t))e_2 \\ &\quad + (W_2 + \Delta W_2(t))e_4 - N g_{ij}^{(1)} \Gamma_1 e_1 - N g_{ij}^{(2)} \Gamma_2 e_3 - \tau_1 N g_{ij}^{(3)} \Gamma_3 e_9 - (\tau(t) - \tau_1) N g_{ij}^{(3)} \Gamma_3 e_{10} - e_{13}] \} \zeta_{ij}(t). \end{aligned} \quad (16)$$

Replacing $\Delta A(t), \Delta W_1(t), \Delta W_2(t), \Delta \Gamma_1(t), \Delta \Gamma_2(t)$, and $\Delta \Gamma_3(t)$ with $HF(t)E_A, HF(t)E_{W_1}, HF(t)E_{W_2}, HF(t)E_{\Gamma_1}, HF(t)E_{\Gamma_2}$ and $HF(t)E_{\Gamma_3}$ respectively, so we have

$$\dot{V}(x_t) \leq \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) [\Pi_0(\tau(t), \dot{\tau}(t)) + \Pi_1^T F(t) \Pi_2 + \Pi_2^T F(t) \Pi_1] \zeta_{ij}(t), \quad (17)$$

where $\Pi_0(\tau(t), \dot{\tau}(t))$, Π_1 and Π_2 are defined in (7). By Lemma 5, it can be deduced that $\varepsilon > 0$ and

$$\dot{V}(x_t) \leq \sum_{1 \leq i < j \leq N} \zeta_{ij}^T(t) [\Pi_0(\tau(t), \dot{\tau}(t)) + \varepsilon^{-1} \Pi_1^T \Pi_1 + \varepsilon \Pi_2^T \Pi_2] \zeta_{ij}(t). \quad (18)$$

Applying the Schur complement Lemma, we know that $\Pi_0(\tau(t), \dot{\tau}(t)) + \varepsilon^{-1} \Pi_1^T \Pi_1 + \varepsilon \Pi_2^T \Pi_2 < 0$, is equivalent to (5). Thus (5) and (6) hold, implies that $\dot{V}(x_t) < 0$, we know that $x^T(t)(U \otimes P)x(t)$ is bounded function and $\lim_{t \rightarrow \infty} \|z_{ij}\| = \lim_{t \rightarrow \infty} \|x_i(t) - x_j(t)\| = 0$. By Lyapunov theorem and Definition 1, the neural networks (4) are globally robustly synchronized. \square

III. MAIN RESULT

In this section, we present a numerical example to illustrate the effectiveness and the reduced conservatism of our result. Consider the dynamical system consisting of model (1), choose the following matrix:

$$A = \begin{bmatrix} 3.5 & 0 \\ 0 & 1.5 \end{bmatrix}, W_1 = \begin{bmatrix} -1 & 0.4 \\ -0.2 & -0.1 \end{bmatrix}, W_2 = \begin{bmatrix} 0.4 & -0.6 \\ 0.2 & 0.4 \end{bmatrix}, \Gamma_1 = \Gamma_2 = \Gamma_3 = \begin{bmatrix} 0.2 & 0 \\ 0 & 0.2 \end{bmatrix}, g^{(1)} = g^{(2)} = g^{(3)} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}, E_A = \begin{bmatrix} 0.2 & 0.1 \\ 0.2 & 0.1 \end{bmatrix},$$

$E_{w_1} = \begin{bmatrix} 0.3 & 0.1 \\ 0.5 & 0.4 \end{bmatrix}, E_{w_2} = \begin{bmatrix} 0.3 & 0.2 \\ 0.1 & 0.3 \end{bmatrix}, E_{\Gamma_1} = E_{\Gamma_2} = E_{\Gamma_3} = \begin{bmatrix} 0.1 & 0 \\ -0.1 & 0.1 \end{bmatrix}, H = \begin{bmatrix} 0.1 & 0 \\ 0 & 0.1 \end{bmatrix}, F(t) = \begin{bmatrix} \sin(t) & 0 \\ 0 & \cos(t) \end{bmatrix}, 0.3 \leq \tau(t) \leq 1,$
 $-0.5 \leq \dot{\tau}(t) \leq 0.5$ and $\sigma = 0.2$. The neural activation functions as follows: $f_1(x_1(t)) = 0.4 \tanh(-x_1), f_2(x_2(t)) = 0.6 \tanh(-x_2)$.
 By applying the MATLAB LMI Toolbox in Theorem1, it yields the following feasible solutions:

$$P = \begin{bmatrix} 0.0347 & 0.0024 \\ 0.0024 & 0.0925 \end{bmatrix}, Q_0 = \begin{bmatrix} 0.0154 & 0.0026 \\ 0.0026 & 0.0647 \end{bmatrix}, S_0 = \begin{bmatrix} 0.0906 & 0.0462 \\ 0.0462 & 0.2326 \end{bmatrix}, S_1 = \begin{bmatrix} 0.3328 & 0.0195 \\ 0.0195 & 0.2772 \end{bmatrix}, S_2 = \begin{bmatrix} 0.1061 & 0.0138 \\ 0.0138 & 0.0875 \end{bmatrix}, S_3 = \begin{bmatrix} 0.2959 & -0.0033 \\ -0.0033 & 0.2527 \end{bmatrix},$$

$$Q_1 = \begin{bmatrix} 0.1459 & 0.0161 & -0.1223 & -0.0068 \\ 0.0161 & 0.3658 & -0.0184 & -0.2557 \\ -0.1223 & -0.0184 & 0.3038 & 0.0011 \\ -0.0068 & -0.2557 & 0.0011 & 0.4078 \end{bmatrix}, Q_2 = \begin{bmatrix} 0.0895 & 0.0027 & 0.0041 & -0.0014 \\ 0.0027 & 0.1914 & -0.0066 & -0.0150 \\ 0.0041 & -0.0066 & 0.1688 & -0.0010 \\ -0.0014 & -0.0150 & -0.0010 & 0.2319 \end{bmatrix}, Q_3 = \begin{bmatrix} 0.0833 & 0.0051 & 0.0124 & -0.0010 \\ 0.0051 & 0.1887 & 0.0009 & 0.0143 \\ 0.0124 & 0.0009 & 0.0925 & 0.0037 \\ -0.0010 & 0.0143 & 0.0037 & 0.1196 \end{bmatrix},$$

$$Y_1 = \begin{bmatrix} 0.0126 & 0.0012 \\ 0.0012 & 0.0384 \end{bmatrix}, Y_2 = \begin{bmatrix} 0.0118 & 0.0022 \\ 0.0022 & 0.0593 \end{bmatrix}, R_1 = \begin{bmatrix} 0.0168 & 0.0009 \\ 0.0009 & 0.0394 \end{bmatrix}, R_2 = \begin{bmatrix} 0.0054 & 0.0028 \\ 0.0028 & 0.0479 \end{bmatrix}, X_1 = \begin{bmatrix} 0.3673 & -0.0111 \\ -0.0111 & 0.2706 \end{bmatrix}, X_2 = \begin{bmatrix} 0.6371 & 0.0048 \\ 0.0048 & 0.8024 \end{bmatrix},$$

$$\Theta_1 = \begin{bmatrix} 0.1091 & 0.0067 \\ 0.0067 & 0.2738 \end{bmatrix}, \Theta_2 = \begin{bmatrix} 0.0238 & 0.0043 \\ 0.0043 & 0.1062 \end{bmatrix}, \Lambda_1 = \begin{bmatrix} 0.4670 & 0.0000 \\ 0.0000 & 0.3147 \end{bmatrix}, \Lambda_2 = \begin{bmatrix} 0.3086 & 0.0000 \\ 0.0000 & 0.3169 \end{bmatrix}, \Lambda_3 = \begin{bmatrix} 0.0748 & 0.0000 \\ 0.0000 & 0.0935 \end{bmatrix}, \quad \varepsilon = 0.1169.$$

The simulation results for the synchronization errors, $z_{ii}(t) = x_i(t) - x_1(t), (i = 2,3)$ of networks (1) are show in Figs.1. The figures show that the networks with the errors converge to zero for given initial values by $x_1(0) = [0.3 \ -0.3]^T, x_2(0) = [0.4 \ -0.35]^T$ and $x_3(0) = [0.1 \ -0.45]^T$.




Fig. 1 Synchronization errors of networks $z_{ii}(t) = x_i(t) - x_1(t), (i = 2,3)$

IV. CONCLUSIONS

In this paper, the robust synchronization analyses of coupled neural networks with interval time-varying delays and leakage delay have been proposed. By constructing a set of improved LKF and Kronecker product properties, sufficient conditions for guaranteeing asymptotic synchronization for the concerned networks have been derived in terms of LMI. A numerical example illustrates the effectiveness of the estimated result.

ACKNOWLEDGMENT

This project was supported by Rajamangala University of Technology Isan.

REFERENCES

1. Gu, K., Kharitonov, V.L. and Chen, J., (2003). Stability of time-delay system, Boston: Birkhauser.
2. Park, M.J., Kwon, O.M., Park, Ju H., Lee, S.M. and Cha, E.J. (2012). Synchronization criteria for coupled neural networks with interval time-varying delays and leakage delay, *Appl. Math. Comput.*, Vol. 218, pp. 6762-6775.
3. Seuret, A. and Gouaisbaut, F. (2013). Wirtinger-based integral inequality: application to time-delay systems. *Automatica*. Vol. 49, pp. 2860-2866.
4. Zhang, X.M. and Han, Q.L. (2014). Global asymptotically stability analysis for delayed neural networks using a matrix-based quadratic convex approach. *Neural Netw.* Vol. 54, pp. 57-69.
5. Du, Y. and Xu, R., (2014). Robust synchronization of an array of neural networks with hybrid coupling and mixed time delays, *ISA Transactions*, Vol. 53, pp. 1015-1023.

6. Cichocki, A. and Unbehauen, R. (1993). *Neural networks for optimization and signal processing*, Wiley, Hoboken, NJ.
7. Kamil, I. A. (2012). Self-synchronization in chaotic systems, *Journal of Engineering and Applied Sciences*, Vol. 7, pp. 411-417.

