

# The External Factors Associated with Stunting Occurrence among 12 – 59 Months Old Toddler

Rico J. Sitorus, Merry Natalia P, Imelda G. Purba, Rini Mutahar, Poppy Fujianti

**Abstract:** *Background: Intervention strategy employed in solving stunting cases should be comprehensive by focusing on both direct (internal) and indirect factors (external). Therefore, the purpose of this study was to observe the indirect factors influencing the occurrence of stunting in toddlers.*

*Methods: The study design used was community-based case control with the population comprising of toddlers between 12-59 months old. The sample size included 49 cases and 98 controls, and the stunting was measured by using anthropometry. Furthermore, short toddlers were indicated by nutrition status which was determined by the ideal height in certain age and compared with the provision of WHO-MGRS (Multicentre Growth Reference Study) that any toddler with z-score less than -2 SD is short and less than -3 SD is categorized as very short.*

*Results: The multivariate analysis conducted proved that toddlers with inadequate child-rearing have a greater risk (Adjusted Odds Ratio (AOR) = 3.03 (95% CI: 1.15-7.98) of experiencing stunting compared to those raised adequately. Those living in families below Guaranteed Minimum Income (GMI) has a greater risk (AOR = 4.63 (95 % CI: 1.701-12.63) of stunting compared to those in families with above Guaranteed Minimum Income (GMI). It was also found that those without exclusive breastfeeding has a greater risk (AOR = 4.53 (95 % CI: 1.64-12.49) compared to those with exclusive breastfeeding.*

*Conclusion: The intervention methods needed to improve stunting's problem-solving in rural areas include giving adequate child-rearing, improving the economic status of the family, and parent's knowledge about giving exclusive breastfeeding for 6 months.*

**Keywords:** *stunting, toddler, rural, breastfeeding*

## I. INTRODUCTION

Stunting is a chronic nutrition problem which is directly caused by the deficiency of macronutrients and micronutrients over a long period. <sup>(1)</sup> Epidemiologically, a health problem occurs because of the imbalance between the host, agent, and the environment, therefore, public health issues like stunting could be attributed to the nutrition deficiency in a host <sup>(2)</sup> as well as external factors such as

**Revised Manuscript Received on September 22, 2019.**

**Rico J. Sitorus**, Epidemiology Department Faculty of Public Health Universitas Sriwijaya, Inderalaya Campus, South Sumatra,30662, Indonesia. rico\_januar@fkm.unsri.ac.id

**Merry Natalia**, Staff in Ministry of Health, Sultan Mahmud Badaruddin II Office, South Sumatra Indonesia.

**Imelda G. Purba**, Environmental Health Department Faculty of Public Health Inderalaya Campus, South Sumatra,30662, Indonesia.

**Rini Mutahar**, Epidemiology Department Faculty of Public Health Universitas Sriwijaya, Inderalaya Campus, South Sumatra,30662, Indonesia.

**Poppy Fujianti**, Environmental Health Department Faculty of Public Health Inderalaya Campus, South Sumatra,30662, Indonesia

access to healthcare service, child-rearing, social economy, poverty, and environment. <sup>(6,7)</sup>

According to the Baseline Health Research of 2013, the prevalence of stunted toddlers in Indonesia reached 37,2%, with very short toddlers being 18% and short toddlers 19,2. <sup>(3)</sup> However, the World Health Organisation (WHO) considers an area to be ideal if the population of the short toddler present there is less than 20%. <sup>(4)</sup>

From the survey conducted by the Nutritional Status Monitoring in 2017, the prevalence of stunting was found to be 29,6% while Nutritional Status Assessment Data in South Sumatera recorded 22,8%. Out of the 17 Districts in South Sumatera, three districts including Musi Rawas Utara, Banyuasin with 32,8% and Ogan Ilir with 29,5% were found to have the highest prevalence. <sup>(5)</sup> Therefore, the reduction of the prevalence of short toddlers is one of the prioritized programs in national development from 2015-2019.

This disease affects the future of a child by leading to various defects such as low intelligence and cognitive ability which would later cause low productivity. Hunt (2005) reported that adult with stunting has lower productivity and income compared to those without. <sup>(8)</sup> Therefore, the aim of this study was to analyze the external factors associated with the occurrence of stunting among 12-59 months old toddlers. The prevalence in Indonesia, as a developing country, is closely related to the population's economic status, access to healthcare intervention, child-rearing, and environmental factor.

## II. METHODS

### A. Study Design and Sampling

The study design used was a community-based case control. The case group consisted of 12-59 months old toddlers with stunting while the control group was made of those without stunting. The information for toddlers with and without stunting was obtained from the Nutritional Status Assessment survey data of 2017 for 147 toddlers.

Stunting was measured by using anthropometry and short toddlers were indicated by nutrition status which was determined by the ideal height in certain age and compared with the provision of WHO-MGRS (Multicentre Growth Reference Study) that any toddler with z-score less than -2 SD is short and less than -3 SD is categorized as very short. <sup>(9)</sup>



## The External Factors Associated with Stunting Occurrence among 12 – 59 Months Old Toddler

Enumerators for anthropometry included people with backgrounds in public health and nutrition, who also received proper training before the conduct of the study. The case group consisted of 49 toddlers while the control group was made of 98 toddlers selected through the use of simple random sampling. Control group were neighbors of case group's toddlers and picked by an unmatching method.

### B. Statistical Analysis

The analysis started with data completion check by editing, coding, and entry. The refined was further analyzed by using the univariate, bivariate and multivariate methods. Univariate analysis was conducted to describe the frequency and distribution of each variable while bivariate analysis described the association of each independent variables to the dependent variable (stunting) which was divided into two categories based on WHO-MGRS (Multicentre Growth Reference Study) standard, stunted and not stunted. Statistic test used in the bivariate analysis was chi-square test while the multivariate analysis made use of multiple logistic regressions test.

### C. Ethical Review

Ethical review for this study was conducted in the Faculty of Public Health, Sriwijaya University.

## III. RESULTS

The proportion of mothers with stunted toddler who have low education level was 20.4%, while those without but with low education level was 48%. The average age of mothers in case and control groups was found to be 32 years old while the stunted toddlers (case) were observed to be 26 months, and those without stunting (control) to be 32 months as shown in Table 1. Toddlers with inadequate child-rearing in the family have a probability 3.03 times greater to become stunted compared to those with adequate child-rearing (Adjusted Odds Ratio (AOR)=3.03;95 % CI 1.15-7.98). Those living in families below Guaranteed Minimum Income (GMI) have a probability 4.63 times greater to become stunted compared to those in families with above Guaranteed Minimum Income (GMI) (Adjusted Odds Ratio (AOR)=4.63; 95 % CI (1.701-12.63). Furthermore, toddlers without exclusive breastfeeding were discovered to have the 4.53 times greater probability to become stunted compared to those with exclusive breastfeeding (Adjusted Odds Ratio(AOR)= 4.53; 95 % CI (1.64-12.49).(Table 3)

**Table 1:** Respondents characteristics

| Variables              | Case |      | Control |      |
|------------------------|------|------|---------|------|
|                        | N    | %    | n       | %    |
| Mother Characteristics |      |      |         |      |
| Education Level        |      |      |         |      |
| Low                    | 10   | 20,4 | 47      | 48   |
| High                   | 39   | 79.6 | 51      | 52   |
| Occupation             |      |      |         |      |
| Unemployed             | 17   | 34.5 | 56      | 57.1 |
| Employed               | 37   | 34.9 | 30      | 61.2 |

|               |                         |                         |
|---------------|-------------------------|-------------------------|
|               | Mean (Min - Max), year  | Mean (Min - Max), year  |
| Mother's age  | 32 (20-45)              | 32 (21-47)              |
|               | Mean (Min - Max), month | Mean (Min - Max), month |
| Toddlers' age | 26 (10-59)              | 32 (11-55)              |

**Table 2:** Independent Factors Associated with Stunting among 12-59 months old toddlers

| Variable                          | Case |      | Contro l |      | p-value | OR,95 % CI          |
|-----------------------------------|------|------|----------|------|---------|---------------------|
|                                   | N    | %    | n        | %    |         |                     |
| Latrine/Toilet Ownership          |      |      |          |      |         |                     |
| Yes                               | 85   | 86.7 | 41       | 83.7 | 0.66    | 0,716 (0,276-1,860) |
| No                                | 13   | 13.3 | 8        | 16.3 |         |                     |
| Family Members                    |      |      |          |      |         |                     |
| ≤ 4 people                        | 11   | 23.4 | 45       | 45.1 | 0.02    | 2.93 (1.35-6.39)    |
| > 4 people                        | 37   | 34.9 | 30       | 61.2 |         |                     |
| Re-occurring diarrhea             |      |      |          |      |         |                     |
| No                                | 70   | 71.4 | 26       | 53.1 | 0,02    | 2.464 (1,214-5,002) |
| Yes                               | 28   | 28.6 | 23       | 46.9 |         |                     |
| Worm Infection                    |      |      |          |      |         |                     |
| Negative                          | 91   | 92.9 | 39       | 20.6 | 0.02    | 3.63 (1,289-10.204) |
| Positive                          | 7    | 7.1  | 10       | 20.4 |         |                     |
| Exclusive Breastfeeding           |      |      |          |      |         |                     |
| Yes                               | 14   | 28.6 | 61       | 62.2 | 0.00    | 4.12 (1.96-8.66)    |
| No                                | 35   | 71.4 | 37       | 37.8 |         |                     |
| Child-rearing                     |      |      |          |      |         |                     |
| Adequate                          | 17   | 34.7 | 56       | 57.1 | 0.01    | 2.51 (1.23-5.11)    |
| Inadequate                        | 34   | 32.1 | 25       | 51   |         |                     |
| Economy Status                    |      |      |          |      |         |                     |
| > Guaranteed Minimum Income (GMI) | 12   | 24.5 | 61       | 62.2 | 0.00    | 5.083 (2.54-10.96)  |
| ≤ Guaranteed Minimum Income (GMI) | 37   | 75.5 | 37       | 37.8 |         |                     |
| Healthcare Facility               |      |      |          |      |         |                     |
| Use                               |      |      |          |      |         | 2.39                |
| Yes                               | 18   | 36.7 | 57       | 51   | 0.02    | (1.182-4.485)       |



No. 31 63. 41 49  
3

**Table 3.** Variables associated with stunting, multivariate analysis

| Variables                      | Case |      | Control |      | Crude OR (95 % CI) | Adjusted OR (95 % CI) |
|--------------------------------|------|------|---------|------|--------------------|-----------------------|
|                                | n    | %    | N       | %    |                    |                       |
| <b>Child-rearing</b>           |      |      |         |      |                    |                       |
| Adequate (reff)                | 17   | 34,7 | 56      | 57.1 | 2.51(1.23-5.11)    | 3.03(1.15-7.98)       |
| Inadequate                     | 32   | 65.3 | 42      | 42.9 |                    |                       |
| <b>Economy Status</b>          |      |      |         |      |                    |                       |
| > UMR (reff)                   | 12   | 24.5 | 61      | 62.2 | 5.083(2.54-10.96)  | 4.63(1.701-12.63)     |
| ≤ UMR                          | 37   | 75.5 | 37      | 37.8 |                    |                       |
| <b>Exclusive Breastfeeding</b> |      |      |         |      |                    |                       |
| Yes (reff)                     | 14   | 28.6 | 61      | 62.2 | 4.12(1.96-8.66)    | 4.53(1.64-12.49)      |
| No                             | 35   | 71.4 | 37      | 37.8 |                    |                       |

#### IV. DISCUSSION

The purpose of this study was to determine the external factors associated with stunting in 12-59 months old toddlers. The results show that family contributes a lot in fulfilling the needs such as nutrition, affection, and child-rearing of toddlers and children. This is possible because child-rearing enables family members to combine knowledge, resources, behaviors, and food safety to promote and maintain a healthy status.<sup>(7-10)</sup> This is consistent with the findings of Reyes H (2004) that the role of the family in child-rearing influences their nutritional status and that there are differences in how it is conducted in rural areas and the cities<sup>(7,11)</sup> However, the nutritional status of children in families can be improved through strategies such as adequate child-rearing, family counselling, home visit by healthcare personnel, and proper training on food and nutrition<sup>(12-14)</sup>.

Furthermore, the economic status of the family was determined by the monthly income. However, those with income below the district regional's standard (Guaranteed Minimum Income) are categorized as a low economy class and vice versa. Moreover, the findings prove that low income is associated with stunting occurrence because families in this class have little access to food and healthcare service when experiencing any form of illness.<sup>(15,16)</sup> Singh (2006) further reported that low economy is associated with poverty which affects nutritional intake because of the

inability to buy foods to be consumed.<sup>(17)</sup> It has also been reported that the wealth of a family has the ability to support the children's nutritional needs, thereby, increasing their growth and cognitive ability maximally.<sup>(18)</sup>

Exclusive breastfeeding is defined to be the intake of nothing other than breastmilk by babies for the first 6 months of their lives except for medicine and vitamins when needed.<sup>(19)</sup> This is important because of the nutritional contents of breastmilk which is needed by babies for their growth. It also contains anti-bacterial and anti-virus that protects them from infections.<sup>(20)</sup> The findings are consistent with the submission of Kuchenbecker et al (2015) that exclusive breastfeeding helps the growth of children below 2 years old.<sup>(21)</sup> It was also reported to have great biological effects and important consequences in achieving nutritional status in children when given for the first six months of birth.<sup>(22)</sup>

#### V. CONCLUSION

The result shows there is a need to pay to some external factors in order to prevent stunting in toddlers in rural areas. They are found to include child-rearing, family's economic status, and exclusive breastfeeding. Therefore, a collaboration between healthcare personnel and the society is required, especially in educating and informing about proper and adequate child-rearing, encouraging exclusive breastfeeding and empowering the society to improve their family's economy.

The limitation of this study enables the possibility of recall and misclassification bias.

#### COMPETING INTERESTS

The Author declares that They have no competing interests.

#### ACKNOWLEDGMENTS

We would like to thank Sriwijaya University as the funder, the Ministry of Health, Puskesmas and all research team members.

This research/article's publication is supported by the United States Agency for International Development (USAID) through the Sustainable Higher Education Research Alliance (SHERA) Program for Universitas Indonesia's Scientific Modeling, Application, Research and Training for City-centered Innovation and Technology (SMART CITY) Project, Grant #AID-497-A-1600004, Sub Grant #IIE- 00000078-UI-1.

#### REFERENCES

- [1] Fikadu T, Assegid S, Dube L. Factors associated with stunting among children of age 24 to 59 months in Meskan district, Gurage Zone, South Ethiopia: a case-control study. BMC Public Health. 2014 Aug;14(1):800.
- [2] Underwood BA. Health and Nutrition in Women, Infants, and Children: Overview of the Global Situation and the Asian Enigma. Nutr Rev. 2002;60.



## The External Factors Associated with Stunting Occurrence among 12 – 59 Months Old Toddler

- [3] NIHRD. Ministry of Health, Report on Basic Health Research 2015. Jakarta: National Institute of Health Research and Development, Ministry of Health Rep of Indonesia; 2015.
- [4] WHO. Childhood Stunting: Context, Causes and Consequences. WHO Conceptual framework. 2013.
- [5] Directorate General of Public Health Ministry of Health, Report Monitoring Nutritional Status, Ministry of Health 2018 Jakarta. (Direktorat Jenderal Kesehatan Masyarakat. Laporan Pemantauan Status Gizi, Kementerian Kesehatan 2018, Jakarta)
- [6] Torlesse H, Cronin AA, Sebayang SK, Nandy R. Determinants of stunting in Indonesian children: evidence from a cross-sectional survey indicate a prominent role for the water, sanitation and hygiene sector in stunting reduction. *BMC Public Health*. 2016 Jul;16(1):669.
- [7] Reyes H, Pérez-Cuevas R, Sandoval A, Castillo R, Santos JI, Doubova S V, et al. The family as a determinant of stunting in children living in conditions of extreme poverty: a case-control study. *BMC Public Health*. 2004 Nov;4(1):57.
- [8] Hunt JM. The potential impact of reducing global malnutrition on poverty reduction and economic development. *Asia Pac J Clin Nutr*. 2005;14.
- [9] WHO. Child growth standards: growth velocity based on weight, length and head circumference: methods and development. World Health Organization; 2009.
- [10] Shinsugi C, Matsumura M, Karama M, Tanaka J, Changoma M, Kaneko S. Factors associated with stunting among children according to the level of food insecurity in the household: a cross-sectional study in a rural community of Southeastern Kenya. *BMC Public Health*. 2015 Apr;15:441.
- [11] Kanjilal B, Mazumdar PG, Mukherjee M, Rahman MH. Nutritional status of children in India: household socio-economic condition as the contextual determinant. *Int J Equity Health*. 2010 Aug;9:19.
- [12] Dewey, K. G. and Adu-Afaruwah S. Systematic review of the efficacy and effectiveness of complementary feeding interventions in developing countries. *itl*. 2008;4:24\*85.
- [13] Roberfroid D, Kolsteren P, Hoeree T, Maire B. Do growth monitoring and promotion programs answer the performance criteria of a screening program? A critical analysis based on a systematic review. *Trop Med Int Heal*. 2005;10(11):1121–33.
- [14] Penny ME, Creed-Kanashiro HM, Robert RC, Narro MR, Caulfield LE, Black RE. Effectiveness of an educational intervention delivered through the health services to improve nutrition in young children: a cluster-randomised controlled trial. *Lancet*. 2005;365(9474):1863–72.
- [15] Delpeuch F, Traissac P, Martin-Prevel Y, Massamba JP, Maire B. Economic crisis and malnutrition: socioeconomic determinants of anthropometric status of preschool children and their mothers in an African urban area. *Public Heal Nutr*. 2000;3.
- [16] Frongillo EA, de Onis M, Hanson KM. Socioeconomic and demographic factors are associated with worldwide patterns of stunting and wasting of children. *J Nutr*. 1997;127.
- [17] Singh MB, Fotedar R, Lakshminarayana J, Anand PK. Studies on the nutritional status of children aged 0–5 years in a drought-affected desert area of western Rajasthan, India. *Public Health Nutr*. 2006/12/01. 2006;9(8):961–7.
- [18] Woldehanna T, Behrman JR, Araya MW. The effect of early childhood stunting on children's cognitive achievements: Evidence from young lives Ethiopia. *Ethiop J Heal Dev = Ya'Ityopya tena lemat mashet*. 2017;31(2):75–84.
- [19] Organization WH. The world health report 2003: shaping the future. World Health Organization; 2003.
- [20] Breastfeeding and the Use of Human Milk. *Pediatrics* [Internet]. 2005 Feb 1;115(2):496 LP-506. Available from: <http://pediatrics.aappublications.org/content/115/2/496.abstract>
- [21] Kuchenbecker J, Jordan I, Reinbott A, Herrmann J, Jeremias T, Kennedy G, et al. Exclusive breastfeeding and its effect on growth of Malawian infants: results from a cross-sectional study. *Paediatr Int Child Health* [Internet]. 2015 Feb;35(1):14–23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25005815>
- [22] Khan MN, Islam MM. Effect of exclusive breastfeeding on selected adverse health and nutritional outcomes: a nationally representative study. *BMC Public Health* [Internet]. 2017 Nov 21;17(1):889. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29162064>

### AUTHORS PROFILE

I am **Rico J. Sitorus**, working with Epidemiology Department Faculty of Public Health Universitas Sriwijaya, Inderalaya Campus, South Sumatra,30662, Indonesia. My area of interest is learning and improving. For contact email me [rico\\_januar@fkm.unsri.ac.id](mailto:rico_januar@fkm.unsri.ac.id)

**My good name is Merry Natalia**, I am Staff in Ministry of Health, Sultan Mahmud Badaruddin II Office, South Sumatra Indonesia.

**Imelda G. Purba**, Environmental Health Department Faculty of Public Health Inderalaya Campus, South Sumatra,30662, Indonesia. My area of interest is learning.

**My good name is Rini Mutahar**, I am working with Epidemiology Department Faculty of Public Health Universitas Sriwijaya, Inderalaya Campus, South Sumatra,30662, Indonesia. My area of interest is public services.

I am **Poppy Fujianti**, currently affiliated with Environmental Health Department Faculty of Public Health Inderalaya Campus, South Sumatra,30662, Indonesia. My area of interest is public health.