

# Urban Sprawl Mapping through Geo-Informatics and Impact on Land use and Landcover of Kakinada Municipal Corporation in Andhra Pradesh, India.

B Sandeep Kumar, R Sivakumar

**Abstract:** *Urban infrastructure and urban sprawl required the idea of preparing a proper management plan to avoid the unwanted environmental and economic impacts that come with it. The main objective of the research is to map the urban sprawl using Geospatial technology and its impact on land use and land cover. The increase in the rate of population over the last two decades is equally responsible for the urban expansion and subsequent infrastructure development. The results of the integrated geospatial study shows that the urban expansion of Kakinada Municipal Corporation was largely caused by the increase in built-up area from 29.67% in 1995, 44.86% in 2011 to 51.34% in 2017 to 62.84% in 2019 out of Kakinada's township area of 189552.6 ha mainly due to natural increase of the population and rural ward migration. Vegetation area was 50.68% in 1995 and has declined to 37.82% in 2011. However, the percentage of vegetation experienced a hike and covered 40.23% in 2017 and then went downhill with a land cover percentage of 34.04% of the total township by the year 2019. Over the last two decades the water-body and the dry land were largely converted into built-up areas. The decline of 49151 ha of water-body due mainly because of the urban expansion and the dry-land lost nearly 27200.79 ha of its land cover to the built-up areas. Therefore, controlling and monitoring of urban expansion using GIS and remote sensing technologies are vital solutions to assess the impact of urban expansion of land use and land cover.*

**Keywords:** *Change detection, Land-use/Land-cover, Geospatial analysis, Urban Sprawl.*

## I. INTRODUCTION

Land-use and land-cover change (LULCC) is a general term for the human modification of Earth's terrestrial surface. The Earth's terrestrial surface modification is monitored on the basis of the LULC (land use and land cover) change. There is a tremendous increase urban population and conversion rural to semi urban and semi urban to urban. Due to this expansion or sprawl of urban leads to the changes in the existing land pattern. This also further reducing the natural resources, changes of many vegetation land and settlement. The urban settlements are growing vertical as well as horizontal. It is very important to understand to rate of change, pattern of change, identification of past land use patten and trend of the change. There is an urgent need to identify and analyze the land use pattern intensified in the past few decades due to the

drastic change in land use and landcover for urban expansion. Hence the aim of this present work was to study and checking urban sprawl redundantly utilizing multirate high, moderate & low-resolution images so that eventually, a procedure is developed where obligation of any harms that occur after some period can be improved on the present field staff in that period. The objectives are in this work is

1. To classify the land use and landcover of Kakinada Municipal corporation of Andhra Pradesh,
2. To map the main LULC classes change trajectories
3. During the study period, in Al Ain city, using remote sensing and GIS techniques.

## II. METHODOLOGY

### A. Study Area


The investigation region is situated in East Godavari district of Andhra Pradesh. The zone shrouded from this examination is around 189 km<sup>2</sup> lying in the middle of 16°52'30''N - 17°6'0''N latitude and 82°4'30''E - 82°24'0''E longitude (Fig.1) The jurisdiction of the corporation is spread over an area of 30.51 km<sup>2</sup> and the population as per the 2011 Census of India was 312,538. It started to expand from 1920, when it was only 20.31 km<sup>2</sup> (7.84 sq. mi) and the population was just over 50,000. The urban agglomeration area of Kakinada is spread over an area of 57.36 km<sup>2</sup> (reference)

**Revised Manuscript Received on June 22, 2020.**

\* Correspondence Author

**B Sandeep Kumar**<sup>1</sup> Department of Civil Engineering, SRM Institute of Science & Technology, Chennai, India. Email: [sandeepkumar\\_sd@srmuniv.edu.in](mailto:sandeepkumar_sd@srmuniv.edu.in)<sup>1</sup>

**R Sivakumar**<sup>2</sup> Department of Civil Engineering, SRM Institute of Science & Technology, Chennai, India. Email: [sivakumr@srmist.edu.in](mailto:sivakumr@srmist.edu.in)<sup>2</sup>



**Fig 1: Study Area**

**B. Climate**

The Kakinada lies over sea flat on 7m. The summers Possess a understanding of precipitation, as the winters possess nothing. Depending on Geiger and Koppen, this setting is assigned. The temperatures in Kakinada is 27.8 °C 82.1 °F. Approximately 11 09 M-M 43.7 inches of all precipitation drops annually.

**C. Transportation Network**

The road network map is prepared 1:50,000 scale toposheets from Survey of India (SOI) and updated using satellite imagery Transportation network of the study area contains mainly Railway with 43.16km and Major Roads covered with 48.04km and National Highway with 47.22km. The legend indicates with colors are red was National Highway and blue was Major Roads and black was railway lane


**D. method**

The methodology involves the analysis of multidade satellite images through Digital image processing and Visual interpretation to prepare various thematic maps like Land use and Land cover, Transportation network etc., A set of Landsat satellite images spanning the years 1995, 2011, 2017 and 2019 was downloaded for Multiyear comparison of land used land cover. This is used as a primary data source for the extraction of the main land-use and land-cover classes. All images are reduced to a common Universal Transverse Mercator (UTM) coordinate system, zone 44 and the WGS84 datum.

**E. Classification**

Supervised classification is used in this paper which led to the following four classed representing most land cover types of the study area:

- Built-up areas
- Vegetation
- Dryland
- Waterbodies.



**Fig 2: Land-use and land cover of year 1995**




Fig 3: Land-use and land cover of year 2011




Fig 5: Land-use and land cover of year 2019




Fig 4: Land-use and land cover of year 2017

### III. RESULTS AND DISCUSSION

#### A. Distribution Of Area




Fig 6: Distribution of area in 1995




Fig 7 : Distribution of area in 2011




Fig 8: Distribution of area in 2017




Fig 9: Distribution of area in 2019

**B. Land use and Land cover Classification**

Land use and land cover data are significant in understanding the surface coverage on the either by natural or manmade. The land spread identifies with the highlights of the earth surface and land use identifies with the human exercises related with a particular zone of land. Change in land spread is customarily estimated utilizing GIS procedures joined with significant ground truthing and measurable examination. There are different remote detecting change discovery techniques that can be utilized to measure changes in land spread. Usually, this includes taking away the phantom reflectance esteems saw in the various groups of pictures caught at various dates that contain coordinating otherworldly and spatial goals. Change in land use, anyway is substantially harder to measure.




Fig 10: Urban Sprawl for 1995, 2011, 2017 and 2019

**C. Urban Sprawl mapping**

Urban areas were extracted from land use and landcover classes of Kakinada municipal corporation using remote sensing and GIS methods (Fig. 1.6) for the period from 1995 to 2019. Urban settlement areas were identified and demarcated the zones of urban based on image processing and interpretation and field knowledge (Table 1.1).

Table 1.1 Urban Expansion for the period between 1995 - 2019

Area in sq.km	1995	2011	2017	2019
Urban area	56.25	85.03	97.33	119.12

**D. Change Detection and analysis**

After the classification of various land use and land cover types using the satellite images and image processing techniques, a comparative analysis was carried out. The compare son was done individually on the basis of the prepared classified land-use and land-cover maps of 1995, 2011, 2017 and 2019. the output of the comparison helps to identify the change in the exiting land use and landcover classes and gives detailed information about the nature of change that occurred during the period between 1995 and 2019.



Further the classified image analysis was used to prepare the change detection or the conversion of the existing LULC to Urban. Hence to get this, Geospatial overlay analysis and spatial metrics are used to identify the changes in the existing urban database. The integrated database on the above all periods were analyses to make the trend of urban expansion. The following Table 1.2 is showing the changes in Built up area, Vegetation, Water body and Dry land for various years mainly 1995, 2011, 2017 and 2019.

The results of the integrated geospatial study show that the urban expansion of Kakinada Municipal Corporation was largely caused by the increase in built-up area from 29.67% in 1995, 44.86% in 2011 increasing to 51.34% in 2017. Further it is expanded to 62.84% in 2019 out of Kakinada’s township area of 189552.6 ha mainly due to natural increase of the population and rural ward migration (Table 1.3).


**Table 1.2 Changes in land use and Land cover and urban expansion (in Sq.km)**

TYPE	1995	2011	2017	2019
Built-up area	56.25	85.03	97.33	119.12
Vegetation	96.06	71.69	76.26	64.53
Waterbody	5.01	3.71	0.79	0.87
Dry land	32.21	29.10	15.16	5.01

Vegetation area was 50.68% in 1995 and has declined to 37.82% in 2011. However, the percentage of vegetation experienced a hike and covered 40.23% in 2017 and then went downhill with a land cover percentage of 34.04% of the total township by the year 2019. Over the last two decades the water-body and the dry land were largely converted into built-up areas. The decline of 49151 ha of water-body due mainly because of the urban expansion and the dry-land lost nearly 27200.79 ha of its land cover to the built-up areas.

**Table 1.3 Changes in land use & Land cover and urban expansion (in %)**

TYPE	1995	2011	2017	2019
Built-up area	29.67	44.86	51.34	62.84
Vegetation	50.68	37.82	40.23	34.04
Waterbody	2.64	1.95	0.41	0.45
Dry land	16.99	15.35	7.99	2.64



**Fig 11: Urban sprawl for the periods between 1995- 2019**

#### IV. CONCLUSIONS

The urbanization and Industrialization are showing a very fast-growing trend for the past few decades. There is an enormous increase of urban population and conversion rural to semi urban and further semi urban to urban. Due to this urban sprawl will lead to the changes in the existing land use and landcover pattern. The natural resources are further declining and also conversion of many agricultural lands to urban settlement. Thus, an exploration study was done on the land use and landcover mapping for in Kakinada Municipal Corporation in East Godavari locale of Andhra Pradesh. The period of observation was planned for about 22 years according to the availability of cloud free satellite data and other relevant data base. There is a huge change in the existing land use and landcover is observed for the time frame between the period 1995 to 2017, The urban settlement territory is expanding constantly according to grouped transient images.

The results of the integrated geospatial study show that the urban expansion of Kakinada Municipal Corporation was largely caused by the increase in built-up area from 29.67% in 1995, 44.86% in 2011 increasing to 51.34% in 2017.

Further it is expanded to 62.84% in 2019 out of Kakinada’s township area of 189552.6 ha mainly due to natural increase of the population and rural ward migration. Vegetation area was 50.68% in 1995 and has declined to 37.82% in 2011

The percentage of vegetation experienced a hike and covered 40.23% in 2017 and then went downhill with a land cover percentage of 34.04% of the total township by the year 2019. Over the last two decades the water-body and the dry land were largely converted into built-up areas.

The decline of 49151 ha of water-body due mainly because of the urban expansion and the dry-land lost nearly 27200.79 ha of its land cover to the built-up areas.

The percentage of vegetation experienced a hike and covered 40.23% in 2017 and then went downhill with a land cover percentage of 34.04% of the total township by the year 2019. Over the last two decades the water-body and the dry land were largely converted into built-up areas. The decline of 49151 ha of water-body due mainly because of the urban expansion and the dry-land lost nearly 27200.79 ha of its land cover to the built-up areas. Hence it requires to have proper planning to safeguard the existing agriculture field and other natural resources. Further, the alarming rate of urban expansion has to be carefully monitored frequently and stop the conversion of fertile lands to urban or semi urban.

#### REFERENCES

- Anderson, J.; Hardy, E.; Roach, J.; Witner, R. A Land Use and Land Cover Classification System for Use with Remote Sensor Data; US Geological Survey Professional Paper 964, USGS: Washington, DC, USA, 1976.
- Abubakar M and Anjide A (2012) Analysis of Land Use/Land Cover Changes to Monitor Urban Sprawl in Keffi-Nigeria. Environmental Research Journal, 6, 130–135.
- Al-Rowaily SL, El-Bana MI, Al-Dujain, FA (2012) Changes in vegetation composition and diversity in relation to morphometry, soil and grazing on a hyper-arid watershed in the central Saudi Arabia. CATENA, 97, 41–49. doi:10.1016/j.catena.2012.05.004

4. Al-Rubaiay TA, Al-Ma'amar AA, Hattab, AA (2010) Integration of remotely sensed data and GIS techniques to study Lesser Zab River Basin. Internalreport No. 3328, Geosurv, Iraq, Baghdad, 73 P. Baghdad.
5. Batty, M., P Longley, P and Fotheringham S 1989, Urban Growth and Form: Scaling, Fractal Geometry, and Diffusion-Limited Aggregation Volume: 21 issue: 11, page(s): 1447- 1472
6. Civco, D.; Hurd, J.; Wilson, E.; Son, M; Zhang, Y. A comparison of land use and land cover change detection methods. In Proceedings of American Society for Photogrammetry and Remote Sensing/American Congress on Surveying and Mapping, Washington, DC, USA, April 2002.
7. Herold, M.; Goldstein, N.; Clarke, K. The spatio-temporal form of urban growth: Measurement, analysis and modeling. Remote Sens. Environ. 2003, 85, 95-105.
8. Lata, M.; Prasad, K.; Bandarinath, K.; Raghavaswamy, R.; Rao, S. Measuring urban sprawl: A case study of Hyderabad. GIS Development 2001, 5. Available online: <http://www.gisdevelopment.net/application/urban/sprawl/urbans0004.htm> (accessed on December 15, 2008).
9. Lakshmi KantaKumar N, Nikhil G Sawant, Shamita Kumar, 2011, Forecasting urban growth based on GIS, RS and SLEUTH model in Pune metropolitan area, International Journal of Geomatics and Geosciences, Volume 2, No 2, 2011, pp: 568-579
10. Mallouk A , Elhadrahi. H, Malaainine M.E. and Rhinane, H, 2019, Using the SLEUTH urban growth model coupled with a GIS to simulate and predict the future urban expansion of Casablanca region, Morocco, The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences, V, Volume XLII-4/W1
11. Moeller, M.S.; Stefanov, W.L.; Netband, M. Characterizing land cover changes in a rapidly growing metropolitan area using long term satellite imagery. In Proceedings of the 2004 Annual Meeting of the American Society for Photogrammetry and Remote Sensing, Denver, CO, USA, 2004.
12. Mohamed Al-shalabi, Lawal Billa, Biswajeet Pradhan, Shattri Mansor and Abubakr A. A. Al-Sharif , 2013 Modelling urban growth evolution and land-use changes using GIS based cellular automata and SLEUTH models: the case of Sana'a metropolitan city, Yemen, Environmental Earth Sciences , 70, pp:425-437 Indian science congress Association
13. Muzein, B.S. Remote sensing and GIS for land cover/land use change detection and analysis in the semi-natural ecosystems and agriculture landscapes of the central Ethiopian Rift Valley. Ph.D. Dissertation, Technische Universität Dresden, Dresden, Germany, 2006
14. Neda Bihamta, Alireza Soffianian, Sima Fakheran and Mehdi Gholamalifard, 2014, Using the SLEUTH Urban Growth Model to Simulate Future Urban Expansion of the Isfahan Metropolitan Area, Iran, Journal of the Indian Society of Remote Sensing volume 43, pages 407-414
15. Qi, Lingrui (2012), Urban land expansion model based on SLEUTH a case study in Dongguan city, China Ph.D. Dissertation, Lund University
16. Sun, H.; Forsythe, W.; Waters, N. Modeling urban land use change and urban sprawl: Calgary, Alberta, Canada. Networks Spatial Econ. 2007, 7, 353-376.
17. Yea-Chung Ding, Yong- Kui Zhang, 2007 , The Simulation Of Urban Growth Applying Sleuth Ca Model To The Yilan Delta In Taiwan, Jurnal Alam Bina, Jilid 09, No: 01, 2007.pp 95-107
18. Yeh, A.; Li, X. Measurement and monitoring of urban sprawl in a rapidly growing region using entropy. Photogramm. Eng. Remote Sensing 2001, 67, 83-90.
19. Yuan, F.; Sawaya, K.; Loeffelholz, B.; Bauer, M. Land cover classification and change analysis of the Twin cities (Minnesota) Metropolitan Area by multitemporal Landsat Remote Sens. Environ. Remote Sens. Environ. 2005, 98, 317-328.



**Mr. R. Sivakumar, Professor**, Professor of Department of Civil Engineering, SRM Institute of Science and Technology, Tamilnadu, India, has graduated B.Sc. in Bharathidasan University, at Tiruchirappalli in 1988 on Geology , received the M.Sc. degree on Geology from Annamalai University in 1990, Tamil Nadu, India and took his M.Tech. on Remote Sensing from Bharathidasan University, Tiruchirappalli, 1993.

His Ph. D degree is on Geoinformatics in Earth Sciences (Catchment Treatment Model) from Bharathidasan University, Tiruchirappalli, 2001, Tamilnadu, India. Member in Indian Society of Remote Sensing. Member in Indian Society of Soil and Water Conservationist and member in Indian Society of Geomatics and Indian Society of Technical Education.

#### AUTHORS PROFILE



**B. Sandeep Kumar** received the B. tech degree in civil engineering from Centurion university of technology and management, Odisha, India in 2014-2018 and doing M. Tech. degree in Remote sensing and GIS from SRM Institute of Science and Technology, Kattankulathur, Chengalpattu District, India in 2018-2020.

